

PhD Thesis

Outline

Sándor Zsuppán

2008

On the Stokes problem

PhD Thesis

Sándor Zsuppán

Thesis supervisor: Dr. Gisbert Stoyan

Professor, Eötvös Loránd University of Sciences, Faculty of Informatics,
Department of Numerical Analysis

Eötvös Loránd University of Sciences, Faculty of Science,
Institute of Mathematics, Doctoral School

Director of the Doctoral School: Dr. Miklós Laczkovich

Professor, Member of the Hungarian Academy of Sciences

Applied mathematics doctoral program

Director of the program: Dr. György Michaletzky

Professor

This thesis was written at

Berzsenyi Dániel Evangélikus Gimnázium (Líceum) és Kollégium

Sopron, 2008

1 Introduction

This thesis is concerned with the Stokes problem, which originates from the Navier-Stokes equations [25] for the motion of a viscous fluid:

$$-\Delta \vec{u} + \text{grad} p = \vec{f} \text{ and } \text{div } \vec{u} = 0. \quad (1)$$

This equation system is to be solved for the unknown velocity \vec{u} and the corresponding pressure p on a planar or spatial domain Ω in appropriate chosen function spaces if one prescribes boundary values for the velocity along the boundary of the domain. The Stokes problem is called of first kind if one prescribes Dirichlet boundary conditions for the velocity.

The thesis contains two main sections.

In the first section we investigate several properties of the Schur complement operator of first kind Stokes problem posed on a plane domain with the help of conformal mapping. Our aim is to show the possible utilization of conformal mapping in reproving already known results and in achieving new ones. We also examine the inf-sup constant which is a number depending only on the shape of the problem domain having a decisive role for the stable solvability of the Stokes problem.

In the second section we give results on the representation of Stokes functions, which are the velocity solutions of the homogeneous Stokes equations (i.e. $\vec{f} = 0$ in (1)) for prescribed Dirichlet boundary conditions. We omit the complex formalism, because we concern also three-dimensional domains. We also generalize these representation formulae to the case of Naviers equation in linear elasticity.

The first section of the thesis is mainly based on my two published papers [53] and [54], while the second section is based on my third paper [55].

2 Results via conformal mapping

In this section we examine mainly the Schur complement operator

$$\mathcal{S} = \text{div } \Delta_0^{-1} \text{grad} : L_2(\Omega) \rightarrow L_2(\Omega), \quad (2)$$

of the first kind Stokes problem, where Δ_0 denotes the vector Laplace operator corresponding to homogeneous Dirichlet boundary values on the boundary of the domain Ω .

We reprove in our implementation the connection between \mathcal{S} and the Friedrichs operator of the domain Ω , which is defined by

$$\mathcal{F} = \mathcal{P} \circ \mathcal{C} : AL_2(\Omega) \rightarrow AL_2(\Omega), \quad (3)$$

where \mathcal{P} is the orthogonal projection of $L_2(\Omega)$ onto the subspace $AL_2(\Omega)$ containing analytic functions, and where \mathcal{C} denotes the conjugacy operator.

This connection, already announced in [13], is formulated in

Theorem 2.1 *Let Ω be a plane domain with enough smooth boundary. Then we have*

$$2\mathcal{S} = \mathcal{I} - \mathcal{C} \circ \mathcal{F} \quad (4)$$

for the Friedrichs operator \mathcal{F} and the Schur complement operator \mathcal{S} of the domain. \square

Remark 2.2 The smoothness of the boundary $\partial\Omega$ of the domain Ω , which plays a decisive role in Theorem 2.1, can be characterized for a simply connected domain with the help of its corresponding Riemann mapping function [37], which maps the domain conformal onto the unit disc D . Theorem 2.1 is also proved for multiply connected domains with enough smooth boundary using a result of [22]. \square

The usage of conformal mapping allows us to give a certain matrix representation for the studied operators (2) and (3). First we observe that

$$\mathcal{F}_D(p) := (\mathcal{F}(f) \circ g)g' \text{ for } p = (f \circ g)g' \in AL_2(D) \quad (5)$$

defines an operator $\mathcal{F}_D : AL_2(D) \rightarrow AL_2(D)$, which is unitary equivalent to \mathcal{F} . The function

$$g : D \rightarrow \Omega \quad (6)$$

denotes here the inverse of the Riemann mapping. The matrix representation comes into scope if we take infinite sequences composed of the Taylor coefficients of the functions in $AL_2(D)$. The operator \mathcal{F}_D is represented then by an infinite matrix \mathcal{M} acting on an appropriate defined space of sequences.

Corollary 2.3 *The Friedrichs operator \mathcal{F} and Schur complement operator \mathcal{S} of the simply connected domain $\Omega = g(D)$ with appropriately smooth conformal mapping g are, respectively, unitary equivalent to the conjugate linear and complex linear operators*

$$\mathcal{M}\mathcal{C} \text{ and } \frac{1}{2}(\mathcal{I} - \mathcal{C}\mathcal{M}\mathcal{C}),$$

which are defined by the infinite matrix \mathcal{M} . \square

Remark 2.4 To be more precise \mathcal{M} acts on the space of complex sequences

$$\ell_{(2,-1)} := \left\{ f \mid f = (f_0, f_1, f_2, \dots)^T \text{ fulfilling } \sum_{n=0}^{\infty} \frac{|f_n|^2}{n+1} < \infty \right\}.$$

The entries of \mathcal{M} are defined by series composed of the coefficients of the Taylor series expansions of the conformal mapping g and of the reciprocal of its derivative. There are domains given by the corresponding function g for which the series defining the entries of \mathcal{M} are divergent. We formulate conditions on the conformal mapping g assuring the existence of the matrix representation. \square

The investigation of \mathcal{F}_D or of the structure of its matrix representant \mathcal{M} reveals several properties of the studied operators in dependence on the problem domain.

Theorem 2.5 *Let η be the conformal map of the domain Ω onto $\tilde{\Omega}$, which are both domains satisfying the conditions of Theorem 2.1. We then have*

$$\|\tilde{\mathcal{F}} - \mathcal{F}\| \leq 2 \sup_{w \in \Omega} |\sin(\arg \eta'(w))|, \quad (7)$$

$$\|\tilde{\mathcal{S}} - \mathcal{S}\| \leq \sup_{w \in \Omega} |\sin(\arg \eta'(w))|. \quad (8)$$

\square

Remark 2.6 The continuity results in Theorem 2.5 can also be formulated in geometric terms involving the boundary of the domain Ω :

$$\|\tilde{\mathcal{F}} - \mathcal{F}\| \leq 2 \max_{w \in \partial\Omega} |\sin(\arg \tilde{\tau}(w) - \arg \tau(w))|, \quad (9)$$

$$\|\tilde{\mathcal{S}} - \mathcal{S}\| \leq \max_{w \in \partial\Omega} |\sin(\arg \tilde{\tau}(w) - \arg \tau(w))|, \quad (10)$$

where $\tau(w)$ denotes the unit tangent vector to $\partial\Omega$, being the conformal image of the unit circle, at the point $w = g(z)$. \square

Lemma 2.7 *Let be $\rho \geq 1$ integer. $\mathcal{M} \in \mathbb{C}^{\rho \times \rho}$ if and only if (6) is a polynomial of order ρ . In this case $\mathcal{M} \in \mathbb{C}^{\rho \times \rho}$ is of rank ρ .* \square

Lemma 2.8 *The matrix \mathcal{M} and the corresponding operators \mathcal{F} and \mathcal{S} are of finite rank if and only if (6) is a fractional rational transformation.* \square

The special class of domains described in Lemma 2.7 are called (classical) quadrature domains and there is extensive research on such domains. The result of Lemma 2.8 is already known for arbitrary quadrature domains, see e.g. [39], where it is proved without the use of conformal mapping. Its use in the proof, however, reveals not only the dimension of the kernel for simply connected domains but also its structure.

Theorem 2.9 *Let Ω be a simply connected quadrature domain of order $\rho + 1$, $\rho \geq 0$. Then the kernel of the associated operators \mathcal{F} and $\mathcal{I} - 2\mathcal{S}$ has codimension $\rho + 1$ and is spanned with the help of the denominator of the Schwarz function of the domain.* \square

Lemma 2.10 *Let $\Omega = g(D)$ have finite area. Then at least one of the entries of \mathcal{M} is not zero. There are domains of infinite area of which corresponding matrix \mathcal{M} is identically the zero matrix.* \square

After the investigation of the operators we study their spectra. We particularly focus on the inf-sup constant $\beta_0(\Omega)$, the square of which is the least positive eigenvalue of the Schur complement operator. We look for possibilities how its eigenvalues can be determined or estimated utilizing the properties of the conformal mapping. Our starting point is an alternate form of the Friedrichs inequality [21]: there exists a constant $\Gamma_\Omega \geq 1$ depending only on the shape of the domain Ω such that for all conjugate harmonic functions u and v in $L_2(\Omega)$ there follows

$$\int_{\Omega} u^2 dA \leq \Gamma_{\Omega} \int_{\Omega} v^2 dA \text{ provided } \int_{\Omega} u v dA = 0. \quad (11)$$

We obtain several results on this basis:

Corollary 2.11 *Let $\Omega = g(D)$ be such that the derivative g' of the conformal map is continuous on the closure of D and $|g'|$ has a positive lower and upper bound on ∂D . Then we have*

$$\frac{1}{\sqrt{2}} \cdot \frac{\inf_{z \in \partial D} |g'(z)|}{\sup_{z \in \partial D} |g'(z)|} \leq \beta_0(\Omega) \leq \frac{1}{\sqrt{2}} \quad (12)$$

for the inf-sup constant of the domain Ω . The equalities hold for $g(z) = z$, that is, $\Omega = D$. \square

Corollary 2.12 *Let the domains Ω and $\tilde{\Omega}$ have smooth boundaries such that the derivative of the bijective conformal mapping η of Ω onto $\tilde{\Omega}$ is continuous on the closure of Ω and $|\eta'|$ has a positive lower and upper bound on $\partial\Omega$. Then there follows*

$$\frac{\inf_{\partial\Omega} |\eta'|}{\sup_{\partial\Omega} |\eta'|} \beta_0(\Omega) \leq \beta_0(\tilde{\Omega}).$$

\square

If η is near to the identity mapping in the sense that the maximum norm of $\eta' - 1$ is small, then we obtain

Theorem 2.13 *If the bijective conformal mapping η of Ω onto $\tilde{\Omega}$ is such that $|\eta'(w) - 1| \leq \varepsilon < 1$ for all $w \in \tilde{\Omega}$, then there follows*

$$\frac{1 - \varepsilon}{1 + \varepsilon} \beta_0(\Omega) \leq \beta_0(\tilde{\Omega}) \leq \frac{1 + \varepsilon}{1 - \varepsilon} \beta_0(\Omega) \quad (13)$$

for the inf-sup constants of the domains. \square

Remark 2.14 Let the domain Ω have $C^{1,\alpha}$ smooth boundary for some $0 < \alpha < 1$. Let the sequence of domains Ω_n , $n = 1, 2, \dots$, tend to the domain Ω in the sense that for their corresponding conformal mappings one has the following: for every $0 < \varepsilon < 1$ there exists a natural number N such that

$$\sup_D |g'_n - g'| < \varepsilon, \quad (14)$$

whenever $n > N$. Then by Theorem 2.13 there follows

$$\lim_{n \rightarrow \infty} \beta_0(\Omega_n) = \beta_0(\Omega),$$

i.e. one has the convergence of the inf-sup constants. Therefore we have obtained a sufficient condition for the convergence of the inf-sup constants of a convergent domain sequence to the inf-sup constant of the limit domain. If one has a condition weaker than (14), then the convergence of the inf-sup constants can not be guaranteed. The example $g_m(z) = z - \frac{c}{m}z^m$, where $m > 1$ is an integer and $0 < |c| \leq 1$ illustrates this phenomenon. If $\Omega_m = g_m(D)$, then the domain sequence (Ω_m) tends to the unit disc but

$$\lim_{m \rightarrow \infty} \beta_0^2(\Omega_m) = \frac{1}{2} - \frac{|c|}{4} < \frac{1}{2} = \beta_0^2(D).$$

We have computed this example with the help of the matrix representation derived for the studied operators. \square

The multiplicity of the eigenvalues of the Schur complement operator can also be examined with the help of conformal mapping.

Theorem 2.15 *Let the bijective conformal mapping of the unit disc be of the form*

$$g(z) = \sum_{n=0}^{\infty} a_{nM+1} z^{nM+1}, \quad (15)$$

where $a_1 \neq 0$ and $M \geq 2$ is an arbitrary integer, then there are eigenvalues of the Schur complement operator with multiplicity more than 1. \square

Remark 2.16 Domains involved in Theorem 2.15 have an M -fold rotational symmetry. We also give an example for a domain such that the square of its inf-sup constant is a double eigenvalue of the Schur complement operator. \square

Remark 2.17 With the help of the matrix representation of the Schur complement operator and of Corollary 2.11 we can estimate the inf-sup constants of some domains (star-shaped domains, convex domains or nearly circular domains for example). We calculate also some examples for domains of which boundary has corners: the wedge, the annular sector, etc. \square

3 Representation results

This section contains results about representation formulae for Stokes functions based mainly on [55]. In this section we omit complex formalism because we also consider three-dimensional domains, but the connection of this section to the previous one is explained.

First we prove the equivalence of two known results from [29] and [30]. This proof is based on a similar representation of conjugate pairs, which are harmonic functions \vec{q} and p on a spatial domain Ω connected by

$$\operatorname{rot} \vec{q}(\vec{x}) = -\nabla p(\vec{x}), \quad \operatorname{div} \vec{q}(\vec{x}) = 0 \quad \text{for } \vec{x} \in \Omega. \quad (16)$$

Lemma 3.1 *Assume $\Omega \subset \mathbb{R}^3$ is a star-shaped domain with respect to the origin. Then the pair (\vec{q}, p) on Ω is a conjugate pair iff there exist scalar harmonic functions φ and ϕ in Ω such that*

$$\vec{q}(\vec{x}) = \nabla \varphi(\vec{x}) + \vec{x} \times \nabla \phi(\vec{x}), \quad (17)$$

$$p(\vec{x}) = \phi(\vec{x}) + \vec{x} \cdot \nabla \phi(\vec{x}), \quad (18)$$

for $\vec{x} \in \Omega$. These harmonic functions are uniquely determined by the pair (\vec{q}, p) under the normalization $\varphi(0) = \phi(0) = p(0) = 0$, and they are given by the formulae

$$\phi(\vec{x}) = \int_0^1 \frac{1}{\tau^2} e^{1-\frac{1}{\tau}} p(\tau \vec{x}) d\tau, \quad (19)$$

$$\varphi(\vec{x}) = \int_0^1 \vec{x} \cdot \vec{q}(\tau \vec{x}) d\tau \left(= \int_0^1 \vec{x} \cdot (\vec{q}(\tau \vec{x}) - \tau \vec{x} \times \nabla \phi(\tau \vec{x})) d\tau \right). \quad (20)$$

\square

Theorem 3.2 *Let $\Omega \subset \mathbb{R}^3$ be a star-shaped domain with respect to the origin. The representation formulae given in [29] and [30] are equivalent.* \square

Remark 3.3 We further connect the results contained in Theorem 3.2 to a third representation – called Papkovitch-Neuber representation – given in [38]. \square

We also study the question how to avoid the restriction of star-shapedness for the domain in Theorem 3.2.

Theorem 3.4 *Let $\Omega \subseteq \mathbb{R}^3$ be a domain. (\vec{v}, p) is a Stokes pair on Ω iff there are harmonic functions \vec{w} and ψ on Ω such that*

$$\vec{v} = -\frac{1}{2}\nabla(\vec{x} \cdot \vec{w}) - \frac{1}{2}\text{rot}(\vec{x} \times \vec{w} + \psi\vec{x}) + \vec{w}, \quad (21)$$

$$p = -2\text{div}\vec{w}. \quad (22)$$

Moreover, these harmonic functions are

$$\vec{w} = \frac{2}{3}\vec{v} - \frac{1}{6}(p\vec{x} - \vec{x} \times \text{rot}\vec{v}), \quad (23)$$

$$\psi = -\frac{1}{6}\vec{x} \cdot \text{rot}\vec{v}. \quad (24)$$

Next we examine similar representations for the solutions Navier's equation for the linear elasticity problem. (This can be seen as another generalization of the formulae given in [29] and [30].)

Theorem 3.5 *Let $\Omega \subset \mathbb{R}^3$ be a star-shaped domain with respect to the origin, and set $\nu \in \mathbb{R}$, $\nu \neq \frac{3}{4}, 1$. The functions $\vec{v} \in C_2(\Omega)$ and $p \in C_1(\Omega)$ satisfy*

$$\Delta\vec{v} = \nabla p, \quad \text{div}\vec{v} = \nu p \quad (25)$$

if and only if there exists a harmonic function $\vec{h} \in C_2(\Omega)$ such that

$$\vec{v}(\vec{x}) = -\frac{1}{2}\left(\vec{x} \text{div}\vec{h}(\vec{x}) + \vec{x} \times \text{rot}\vec{h}(\vec{x})\right) + \left(\frac{3}{2} - 2\nu\right)\vec{h}(\vec{x}), \quad (26)$$

$$p(\vec{x}) = -2\text{div}\vec{h}(\vec{x}), \text{ for } \vec{x} \in \Omega. \quad (27)$$

The harmonic function \vec{h} is unique, and we have

$$\vec{h}(\vec{x}) = \frac{2}{3-4\nu}\left(\vec{v}(\vec{x}) - \frac{1}{4}\left(p(\vec{x})\vec{x} - \frac{1}{1-\nu}\vec{x} \times \text{rot}\vec{v}(\vec{x})\right) + \vec{x} \times \nabla\phi(\vec{x})\right), \quad (28)$$

where the function ϕ is harmonic in Ω and defined by

$$\phi(\vec{x}) = -\frac{1}{4(1-\nu)}\int_0^1 t^{4(1-\nu)}\vec{x} \cdot \text{rot}\vec{v}(t\vec{x})dt. \quad (29)$$

\square

To complete the generalization of the representation theorems we consider also the two-dimensional counterpart of Theorem 3.5.

Theorem 3.6 *Let $\Omega \subseteq \mathbb{R}^2$ be a domain and set $\nu \in \mathbb{R}$, $\nu \neq \frac{1}{2}, 1$. The functions $\vec{v} \in C_2(\Omega)$ and $p \in C_1(\Omega)$ satisfy (25) iff there exists a harmonic function $\vec{h} \in C_2(\Omega)$ such that*

$$\vec{v}(\vec{x}) = -\frac{1}{2} \left(\vec{x} \operatorname{div} \vec{h}(\vec{x}) + \vec{x}^\perp \operatorname{rot} \vec{h}(\vec{x}) \right) + (1 - 2\nu) \vec{h}(\vec{x}) \quad (30)$$

$$p(\vec{x}) = -2 \operatorname{div} \vec{h}(\vec{x}) \text{ for } \vec{x} \in \Omega; \quad (31)$$

this harmonic function \vec{h} is unique, and we have

$$\vec{h}(\vec{x}) = \frac{1}{1 - 2\nu} \left(\vec{v}(\vec{x}) - \frac{1}{4} \left(p(\vec{x}) \vec{x} - \frac{1}{1 - \nu} \vec{x}^\perp \operatorname{rot} \vec{v}(\vec{x}) \right) \right) \text{ for } \vec{x} \in \Omega. \quad (32)$$

□

Remark 3.7 The formulae (30) and (31) can be put in complex form equivalent to the formulae used in the previous Section 2. □

We end this section by formulating two connections between the two-dimensional and three-dimensional representation formulae.

Acknowledgement

I would like to thank my supervisor Prof. Gisbert Stoyan for suggesting the subject of this thesis and also for his continuous support and encouragement during my work.

References

- [1] Adams R.A., Sobolev Spaces, Pure and Appl. Math. 65., Academic Press, New York (1975)
- [2] Aharonov D., Shapiro H.S., Domains in which analytic functions satisfy quadrature identities, J. Analyse Math. 30 (1976), 39–73.
- [3] Aristov P.P. and Chizhonkov E.V., On the constant in the LBB condition for rectangular domains. Report No. 9534, Dept. Math., Univ. of Nijmegen, Sept. 1995.

- [4] Arushanyan I.O. and Chizhonkov E.V., Numerical investigation of the constant in the inf-sup inequality using boundary integral equations. In: Packages of Programmes for Applications, V.A. Morozov, O.B. Arushanyan (eds.), Editing House of Moscow Univ., 1997, 49–59.
- [5] Babuška I., The finite element method with Lagrangian multipliers, Numer. Math. 20. (1973), 179–192.
- [6] Bergman S., Schiffer M., Kernel functions and conformal mapping, Composition Math. 8. (1951), 205–249.
- [7] Bourdon P.S., Density of the polynomials in Bergman spaces, Pacific Journal of Mathematics 130, No. 2 (1987), 215–221.
- [8] Brezzi F., On the existence, uniqueness and approximation of saddle-point problems arising from Lagrange multipliers, R.A.I.R.O. Anal. Numer. 8. (1974), 129–151.
- [9] Brezzi F., Fortin M., Mixed and Hybrid Finite Element Methods, Springer, New York, (1991).
- [10] Chizhonkov E.V., On the constant in the LBB condition for ring domains. Report No. 9537, Dept. Math., Univ. of Nijmegen, Sept. 1995.
- [11] Chizhonkov E.G., On spectral properties of the pressure operator induced by the Stokes equations. In: Proceedings of the Conference in memory of the 175th birthday of P.L. Chebyshev, v. 2. Editing House of the Mech.-Math. Faculty of Moscow Univ., 1996, 363–366.
- [12] Chizhonkov E.V. and Olshanskii M.A., On the domain geometry dependence of the LBB condition. Math. Modell. Numer. Analysis 34, 5 (2000), 935–951.
- [13] Costabel M. and Dauge M.: On the Cosserat spectrum in polygons and polyhedra, IRMAR Conference Lausanne 2000.
- [14] Crouzeix M., Étude d’une méthode de linéarisation. Résolution numérique des équations Stokes stationnaires. In: Cahier de l’INRIA 12, 1974, 139–244.
- [15] Crouzeix M., On an operator related to the convergence of Uzawa’s algorithm for the Stokes equation. In: Computational Science for the 21st Century (J. Périaux et al., eds.). New York: Wiley 1997, 242–249.

- [16] Dobrowolski M., On the LBB constant on stretched domains, *Mat. Nachr.*(2003), 254–255, 64–67.
- [17] Dobrowolski M., On the LBB condition in the numerical analysis of the Stokes equations, *Appl. Numer. Math.* 54, No. 3-4 (2005), 314–323.
- [18] Dobrowolski M., Stoyan G., Algebraic and discrete Veltre decompositions, *BIT* 41(3). (2001), 465–479.
- [19] Duren P.L., Theory of H^p spaces, *Pure and Appl. Math.* 38., Academic Press, New York (1970)
- [20] Duren P.L., *Univalent Functions*, Springer-Verlag, New York (1983)
- [21] Friedrichs K., On certain inequalities and characteristic value problems for analytic functions and for functions of two variables. *Trans. AMS* 41 (1937), 321–364.
- [22] Garabedian P.R., A partial differential equation arising in conformal mapping, *Pacific J. Math.* 1(1951), 485–524.
- [23] Garcia S.R., Means of unitaries, conjugations, and the Friedrichs operator, *J. Math. Anal. Appl.* 335 (2007), 941–947.
- [24] Gaultier M., Lezaun M., Quelques propriétés du spectre de l'opérateur $-\operatorname{div}(-\Delta)^{(-1)}\nabla$ dans L^2 , *C.R. Acad. Sci.(Paris)* 315 (1992), 321–364.
- [25] Girault V., Raviart P.A., *Finite Element Methods for Navier-Stokes Equations*, Springer, New York (1986)
- [26] Horgan C.O. and Payne L.E., On inequalities of Korn, Friedrichs and Babuška-Aziz. *Archive Rat. Mech. Anal.* 82 (1983), 165–179.
- [27] Jakobsson S., The harmonic Bergman kernel and the Friedrichs operator, *Ark. Mat.*, 40 (2002), 89–104.
- [28] Kellogg O.D., On the derivatives of harmonic functions on the boundary, *Trans. Amer. Math. Soc.* 33(2)(1931), 486–510.
- [29] Kratz W., On the representation of Stokes flows, *SIAM J. Math. Anal.* 22(1991)2, 414–423.
- [30] Kratz W., Lindae A., A representation formula for three-dimensional Stokes flows, *Zeitschrift für Analysis und ihre Anwendungen* 11(1992)3, 371–375.

- [31] Kratz W. and Peyerimhoff A., A numerical algorithm for the Stokes problem based on an integral equation for the pressure via conformal mappings, *Numerische Mathematik* 58 (1990), 255–272.
- [32] Ladyzhenskaya O.A., *The Mathematical Theory of Viscous Incompressible Flow*, Gordon and Breach, New York (1969).
- [33] Lin P., Rochberg R., On the Friedrichs operator, *Proc. Amer. Math. Soc.* 123 (1995), 3335–3342.
- [34] Maday Y., Meiron D., Patera A.T., Ronquist E.M., Analysis of iterative methods for the steady and unsteady Stokes problem: application to spectral element discretizations, *SIAM J. Sci. Comput.* 14 (1993), 310–337.
- [35] Mikhlin S.G., The spectrum of the operator pencil of elasticity theory, *Uspekhi Mat. Nauk* 28, No. 3 (171), (1973), 43–82.
- [36] Nečas J., *Equations aux Dérivées Partielles*, Presses de l'Université de Montréal, (1965).
- [37] Nehari Z., *Conformal Mapping*. McGraw-Hill Book Company, Inc., New York, 1952.
- [38] Neuber H., Ein neuer Ansatz zur Lösung räumlicher Probleme der Elastizitätstheorie, *Z. Angew. Math. Mech.* 14(1934), 203–212. (German)
- [39] Putinar M., Shapiro, H.S., The Friedrichs operator of a planar domain, *Oper. Theory Adv. Appl.*, 113, Birkhäuser Verlag, Basel (2000), 303–330.
- [40] Putinar M., Shapiro, H.S., The Friedrichs operator of a planar domain II, *Oper. Theory Adv. Appl.*, 127, Birkhäuser Verlag, Basel (2001), 519–551.
- [41] Razali M. R. M., Nashed M. Z. and Murid A. H. M., Numerical conformal mapping via the Bergman kernel, *Journal of Computational and Applied Mathematics* 82 (1997), 333–350.
- [42] Sakai M., Null quadrature domains, *J. Analyse Math* 40 (1981), 144–154.
- [43] Sakai M., *Quadrature domains*, Lecture Notes in Math. 934, Springer-Verlag (1982), 287–331.

- [44] Shapiro H.S., Unbounded quadrature domains, Lecture Notes in Math., Springer-Verlag (1987), 287–331.
- [45] Stoyan G., Towards discrete Velte decompositions and narrow bounds for inf-sup constants, Computers & Maths. with Appls., 38, 7-8, (1999), 243–261.
- [46] Stoyan G., Iterative Stokes solvers in the harmonic Velte subspace, Computing 67 (2000), 13–33.
- [47] Stoyan G., $-\Delta = -\text{grad div} + \text{rot rot}$ for matrices, with application to the finite element solution of the Stokes problem, East-West J. Numer. Math. 8., (2000) 323–340
- [48] Stoyan G., Strauber Gy., Baran Á., Generalizations to discrete and analytical Crouzeix-Velte decompositions, Numerical Linear Algebra, 11 (2004), 565–590.
- [49] Velte W., On optimal constants in some inequalities. In: The Navier-Stokes Equations, Theory and Numerical Methods (Heywood J.G. et al., eds.). Lecture Notes in Math. 1431. Springer-Verlag Berlin 1990, 158–168.
- [50] Velte W., On inequalities of Friedrichs and Babuška-Aziz, Meccanica 31 (1996), 589–596.
- [51] Warschawski S., Über das Randverhalten der Ableitung der Abbildungsfunktion bei konformer Abbildung (in german), Math. Zeitschrift 35 (1932), 21–456.
- [52] Zimmer S., Rand-Druckkorrektur für die Stokes-Gleichung, Thesis, Techn. Univ. München, 1996.
- [53] Zsuppan S., On the spectrum of the Schur complement of the Stokes operator via conformal mapping, Methods and Applications of Analysis, Vol.11, No.1 (2004), 133–154.
- [54] Zsuppan S., On connections between the Stokes-Schur and the Friedrichs operator, with applications to the inf-sup problem, Annales Univ. Sci. Budapest., 48.(2005), 151–171.
- [55] Zsuppan S., On representations of Stokes flows and of the solutions of Naviers equation for linear elasticity, Analysis, Vol. 28, Issue 2 (2008), 219–237.